

# Guideline for the treatment of no light perception eyes induced by mechanical ocular trauma

Hua Yan<sup>1,2</sup>  | Kehu Yang<sup>3,4</sup>  | Zhizhong Ma<sup>5</sup> | Ferenc Kuhn<sup>6</sup> | Wenfang Zhang<sup>7</sup> | ZhiJun Wang<sup>8</sup> | Yuntao Hu<sup>9</sup> | Hai Lu<sup>10</sup> | Yoshida Shigeo<sup>11</sup> | Gungor Sobaci<sup>12</sup> | Sengul Ozdek<sup>13</sup> | Matteo Forlini<sup>14,15</sup> | Bo Huang<sup>16</sup> | Yannian Hui<sup>17</sup> | Ming Zhang<sup>18</sup> | Gezhi Xu<sup>19</sup> | Wenbin Wei<sup>10</sup> | Yanrong Jiang<sup>20</sup> | DongHo Park<sup>21</sup> | RodrigoAntonio Brant Fernandes<sup>22</sup> | Yuguang He<sup>23</sup> | Andres Rousselot<sup>24</sup>  | Annette Hoskin<sup>25</sup> | Gangadhara Sundar<sup>26</sup> | Yong Liu<sup>27</sup> | Yusheng Wang<sup>17</sup> | Lijun Shen<sup>28</sup> | Haoyu Chen<sup>29</sup> | Huijin Chen<sup>5</sup> | Gezhi Han<sup>30</sup> | Rui Jiang<sup>19</sup> | Xuemin Jin<sup>31</sup> | Jijian Lin<sup>32</sup> | Jing Luo<sup>33</sup> | Zhaoyang Wang<sup>34</sup> | Yong Wei<sup>28</sup> | Ying Wen<sup>35</sup> | Zhenggao Xie<sup>36</sup> | Yi Wang<sup>37</sup> | Xun Yang<sup>38</sup> | Wenzhen Yu<sup>20</sup> | Zhi Zheng<sup>39</sup> | Xiaodong Sun<sup>39</sup> | Jianhong Liang<sup>20</sup> | Qin Liu<sup>40</sup> | Jinguo Yu<sup>1</sup> | Shihui Wei<sup>41</sup>  | Zhengxiang Li<sup>1</sup> | Lu Chen<sup>42</sup> | Xiufen Wang<sup>1</sup> | Lili Wei<sup>3</sup> | Haokun Zhang<sup>1</sup> | Siyue Chen<sup>1</sup> | Yuming Liu<sup>1</sup> | Xu Guo<sup>1</sup> | Siyuan Liu<sup>7</sup> | Xinhua Xu<sup>40</sup> | Yibo Tao<sup>7</sup> | Yixuan Chen<sup>40</sup> | Yaolong Chen<sup>3,4</sup> 

<sup>1</sup>Department of Ophthalmology, Tianjin Medical University General Hospital, Tianjin, P. R. China

<sup>2</sup>School of Medicine, Nankai University, Tianjin, P. R. China

<sup>3</sup>Evidence-based Medicine Center, School of Basic Medical Sciences, Lanzhou University, Lanzhou, P. R. China

<sup>4</sup>WHO Collaborating Centre for Guideline Implementation and Knowledge Translation, Lanzhou University, Lanzhou, P. R. China

<sup>5</sup>Peking University Eye Center, Peking University Third Hospital, Beijing, P. R. China

<sup>6</sup>Department of Ophthalmology Research, Helen Keller Foundation for Research and Education, Birmingham, Alabama, USA

<sup>7</sup>Department of Ophthalmology, Lanzhou University Second Hospital, Lanzhou, P.R. China

<sup>8</sup>Department of Ophthalmology, China-Japan Friendship Hospital, Beijing, P. R. China

<sup>9</sup>Department of Ophthalmology, Beijing Tsinghua Changgung Hospital, School of Clinical Medicine, Tsinghua University, Beijing, P. R. China

<sup>10</sup>Beijing Ophthalmology and Visual Science Key Laboratory, Beijing Tongren Eye Center, Beijing Tongren Hospital, Capital Medical University, Beijing, P. R. China

<sup>11</sup>Department of Ophthalmology, Kurume University School of Medicine, Kurume, Fukuoka, Japan

<sup>12</sup>Department of Ophthalmology, Faculty of Medicine, Hacettepe University, Ankara, Turkey

<sup>13</sup>Department of Ophthalmology, Gazi University Faculty of Medicine, Ankara, Turkey

<sup>14</sup>Department of Ophthalmology, Domus Nova Hospital, Ravenna, Italy

<sup>15</sup>Department of Ophthalmology, Ospedale dello Stato della Repubblica di San Marino, Ravenna, Italy

<sup>16</sup>Department of Ophthalmology, University of Mississippi Medical Center, Jackson, Mississippi, USA

<sup>17</sup>Department of Ophthalmology, Xijing Hospital, Eye Institute of Chinese PLA, Fourth Military Medical University, Xi'an, P. R. China

<sup>18</sup>Department of Ophthalmology, Sichuan University West China Hospital, Chengdu, P. R. China

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2022 The Authors. *Journal of Evidence-Based Medicine* published by Chinese Cochrane Center, West China Hospital of Sichuan University and John Wiley & Sons Australia, Ltd.

- <sup>19</sup>Eye Institute and Department of Ophthalmology, Eye and ENT Hospital, Fudan University, Shanghai, P. R. China
- <sup>20</sup>Department of Ophthalmology, Peking University People's Hospital, Eye Diseases and Optometry Institute, Beijing, P. R. China
- <sup>21</sup>Department of Ophthalmology, School of Medicine, Kyungpook National University, Kyungpook National University Hospital, Daegu, Republic of Korea
- <sup>22</sup>Department of Ophthalmology and Visual Sciences, Escola Paulista de Medicina, Federal University of Sao Paulo, Sao Paulo, Brazil
- <sup>23</sup>Department of Ophthalmology, UT Southwestern Medical Center, Dallas, Texas, USA
- <sup>24</sup>Department of Ophthalmology, Universidad del Salvador, Buenos Aires, Argentina
- <sup>25</sup>Save Sight Institute, The University of Sydney, Sydney, New South Wales, Australia
- <sup>26</sup>Department of Ophthalmology, National University Hospital of Singapore, Singapore, Singapore
- <sup>27</sup>Department of Ophthalmology, First Affiliated Hospital of Army Medical University of PLA, Chongqing, P. R. China
- <sup>28</sup>School of Ophthalmology and Eye Hospital, Wenzhou Medical University, Wenzhou, P. R. China
- <sup>29</sup>Joint Shantou International Eye Center, Shantou University and the Chinese University of Hong Kong, Shantou, P. R. China
- <sup>30</sup>Clinical College of Ophthalmology, Tianjin Medical University, Tianjin Key Lab of Ophthalmology and Visual Science, Tianjin Eye Institute, Tianjin Eye Hospital, Tianjin, P. R. China
- <sup>31</sup>Department of Ophthalmology, The First Affiliated Hospital of Zhengzhou University, Henan Province Eye Hospital, Henan International Joint Research Laboratory for Ocular Immunology and Retinal Injury Repair, Zhengzhou, P. R. China
- <sup>32</sup>Eye Center, Second Affiliated Hospital, School of Medicine, Zhejiang University, Hangzhou, P. R. China
- <sup>33</sup>Department of Ophthalmology, The Second Xiangya Hospital of Central South University, Changsha, P. R. China
- <sup>34</sup>Department of Ophthalmology, The Shanghai Tenth People's Hospital of Tongji University, Shanghai, P. R. China
- <sup>35</sup>Department of Ophthalmology, Affiliated Eye Hospital of Shandong University of Traditional Chinese Medicine, Jinan, P. R. China
- <sup>36</sup>Department of Ophthalmology, Nanjing Drum Tower Hospital, the Affiliated Hospital of Nanjing University Medical School, Nanjing, P. R. China
- <sup>37</sup>Chongqing Aier General Hospital, Aier School of Ophthalmology, Central South University, Chongqing, P. R. China
- <sup>38</sup>Department of Fundus Diseases and Ocular Trauma, LiXiang Eye Hospital of Soochow University, Suzhou, P. R. China
- <sup>39</sup>Department of Ophthalmology, Shanghai General Hospital, Shanghai, P. R. China
- <sup>40</sup>Department of Ophthalmology, Gansu Provincial Hospital, Lanzhou, P. R. China
- <sup>41</sup>Department of Neuro-Ophthalmology, Chinese People's Liberation Army General Hospital, Beijing, P. R. China
- <sup>42</sup>School of Finance, Nankai University, Tianjin, P. R. China

## Correspondence

Hua Yan, Department of Ophthalmology,  
Tianjin Medical University General Hospital,  
Tianjin, 300052, China.  
Email: [zyyyanhua@tmu.edu.cn](mailto:zyyyanhua@tmu.edu.cn)

Kehu Yang, Evidence-based Medicine Center  
of Lanzhou University/WHO Collaborating  
Centre for Guideline Implementation and  
Knowledge Translation, Lanzhou, 730000,  
China.  
Email: [yangkehu-ebm@lzu.edu.cn](mailto:yangkehu-ebm@lzu.edu.cn)

## Abstract

Severe mechanical ocular trauma with no light perception (NLP) predicts a poor prognosis of visual acuity and enucleation of the eyeball. Since the innovative treatment concept of exploratory vitreoretinal surgery has developed and treatment technology has advanced, the outcomes of severe ocular trauma treatment in NLP patients have greatly improved. However, there remains a lack of unified standards for the determination, surgical indication, and timing of vitrectomy in NLP eye treatment. To address these problems, we aimed to create evidence-based medical guidelines for the diagnosis, treatment, and prognosis of mechanical ocular trauma with NLP. Sixteen relevant recommendations for mechanical ocular trauma with NLP were obtained, and a consensus was reached. Each recommendation was explained in detail to guide the treatment of mechanical ocular trauma associated with NLP.

## KEYWORDS

evidence-based medicine, mechanical ocular trauma, no light perception, treatment guidelines

## 1 | INTRODUCTION

The concept of no light perception (NLP) after mechanical ocular trauma was first introduced by Pieramici in 1997.<sup>1</sup> There is a lack

of epidemiological data on NLP eyes induced by mechanical ocular trauma, and the geographical specificity is obvious in the existing epidemiological studies on mechanical ocular trauma. The proportion of NLP eyes was 6%–22% in severe open globe injury.<sup>2–4</sup> NLP

eyes induced by mechanical ocular trauma generally have serious eye injuries.<sup>5</sup> In the past, surgical treatment was generally considered meaningless when the visual function was severely damaged to NLP after ocular trauma, and enucleation might be inevitable.<sup>1,6</sup> Nevertheless, with improvements in vitreoretinal surgical techniques and cumulative experiences in ocular trauma treatment, the visual acuity of patients with NLP eyes may partially improve.<sup>7</sup> Hence, mechanical ocular trauma with an initial NLP is not an indication of permanent vision loss or enucleation.<sup>8,9</sup> However, there is still a lack of unified standards for the identification of NLP eyes, determination of surgical indications, principles, and timing of surgery in NLP eyes.

Ophthalmologists often make diagnoses and treatments based on their experience because of the complexity and variability of mechanical eye trauma. Evidence-based medicine methods are urgently needed to formulate high-quality clinical guidelines to standardize the treatment of NLP eyes induced by mechanical ocular trauma. Therefore, worldwide ophthalmologists in the field of ocular trauma, in conjunction with the World Health Organization (WHO) Collaborating Centre for Guideline Implementation and Knowledge Translation, have developed the "Guideline for the treatment of no light perception eyes induced by mechanical ocular trauma, 2022" using the methods and procedures of evidence-based guidelines, hoping to standardize the treatment of NLP eyes induced by mechanical ocular trauma.

## 2 | METHODS

The guideline is based on the WHO guidebook published in 2014,<sup>10</sup> referring to the definition of clinical practice guidelines proposed by the American Academy of Medical Sciences<sup>11</sup> and following the Appraisal of Guidelines for Research and Evaluation II<sup>12</sup> and Reporting Items for Practice Guideline in Health Care.<sup>13</sup>

### 2.1 | Origination, development and support units for the guideline

This guideline was developed by Tianjin Medical University General Hospital, and methodological support was provided by the Grading of Recommendations Assessment, Development, and Evaluation (GRADE) China Center/Evidence-based Medicine Center of Lanzhou University. The guideline was registered on the International Practice Guideline Registration Platform (<http://www.guideline-registry.cn>), registration number: IPGRP-2021CN008.

### 2.2 | Guideline users and target groups

This guideline is applicable to all medical institutions that manage the treatment of mechanical ocular trauma with NLP. The guideline users are ophthalmologists, and the primary target population is patients diagnosed with NLP induced by mechanical ocular trauma.

### 2.3 | Working group on guideline development steering

The guideline development working group was composed of the Committee of Editorial Group, Consensus Expert Group, Guideline Development Group, and Audit Expert Group. The Committee of the Editorial Group and Consensus Expert Group consisted of 53 experts from four disciplines/specialties from 10 countries, patient representatives participating in the guideline formulation process, and legal advisers participating in the external review of the guideline.

### 2.4 | Clinical problem selection

The research group systematically searched for guidelines, systematic evaluations, and original studies related to the treatment of mechanical ocular trauma with NLP. Combined with the results from in-depth interviews with stakeholders, a list of clinical problems and outcome indicators of the guideline was initially prepared and then classified and merged. A total of 60 ophthalmologists were surveyed using a questionnaire. We used the Delphi method to select and determine the 16 clinical problems from this guideline. Clinical experts and scientists developed clinical problems based on the Population, Intervention, Comparison, and Outcome principle. The outcome indicator list was proposed based on the literature research, in-depth interviews, and patients' wishes and values. Finally, the Guideline Development Working Group finalized the outcome indicators of this guideline after several discussions.

### 2.5 | Evidence collection

Relevant English literature was queried through multiple databases such as PubMed, Embase, The Cochrane Library, and Web of Science. Chinese literature was queried using China National Knowledge Infrastructure (CNKI), China Biology Medicine disc (CBM), WanFang Data, and VIP databases. Common clinical guideline websites included the National Guideline Library, International Guideline Collaboration Network, Inter-School Guideline Network in Scotland, National Institute for Clinical Excellence, and WHO. The retrieval period was from the database construction to June 2021. References in the included literature were thoroughly studied.

### 2.6 | Evidence screening and extraction

Research evidence included existing clinical guidelines, systematic reviews, randomized controlled trials, cohort studies, case-control studies, and cross-sectional studies, which included the diagnosis and treatment of mechanical ocular trauma with NLP. The literature was graded by reading the titles, abstracts, and full texts. The information included in the study was extracted according to

**TABLE 1** Grading of evidence quality and recommendation strength

Grading of evidence quality	Detailed description
High (A)	Having high certainty on that the observation is close to the real value
Medium (B)	Having moderate certainty on observations: Observations may be close to the real value, but they may vary greatly
Low (C)	Having limited certainty on observations: Observations can be very different from real values
Very Low (D)	Having little certainty on observations: They can be extremely different from real values
Grading of recommendation strength	Detailed description
Strong (1)	It is clearly shown that interventions do more benefit or harm
Weak (2)	The advantages and disadvantages are uncertain or the evidence, regardless of quality, shows that the advantages and disadvantages are equal

a predesigned data extraction table. The screening and information extraction of each paper were carried out independently by two professionals, and a third party was involved if there were differences of opinions.<sup>14,15</sup>

## 2.7 | Evaluation and grading of evidence quality

AMSTAR-2<sup>16</sup> was used to evaluate the methodological quality of the included system. Evaluation/meta-analysis, and the Cochrane bias risk assessment tool, QUADAS-2, and Newcastle-Ottawa-Scale (NOS) were used to evaluate the methodological quality of the corresponding clinical trials. The screening and information extraction of each paper were carried out independently by two professionals, and a third party was involved if there were differences of opinions.<sup>17–19</sup>

GRADE<sup>20,21</sup> was adopted to evaluate the quality of evidence and strength of recommendation for each clinical issue. A good practice statement (GPS) was used to evaluate the qualitative research.<sup>22</sup> Table 1 shows the grading of the evidence quality and recommendation strength.

## 2.8 | Patient preferences and values

A questionnaire on clinical issues that catered to diverse patient preferences and values was developed by the Guideline Development Working Group. Patients with mechanical ocular trauma ( $n = 100$ ) were selected to complete this questionnaire. The findings were ana-

lyzed statistically, collated by members of the guideline development team, and considered in the formulation of recommendations.

## 2.9 | Development and updating of recommendations

The Guideline Development Working Group developed 16 recommendations based on the evidence of meta-analysis and other factors, including the patient's values and personal preferences, the cost of interventions, and balancing the pros and cons. All recommendations reached consensus through a one-round Delphi questionnaire.

## 2.10 | Audit and approval

After reaching consensual recommendations reached, a draft of the guideline was submitted to the Audit Expert Group for review, which comprised ocular trauma experts, guideline formulation methodology experts, legal advisers, and patient representatives. The Guideline Development Working Group revised the overall opinion based on feedback from the Audit Expert Group. Finally, the Consensus Expert Group discussed and approved the guideline.

## 2.11 | Dissemination and implementation of the guideline

Upon the release of the guideline, the Guideline Development Working Group will disseminate, implement, and promote it in the following ways: (1) Interpret them at relevant academic conferences, (2) publish them in academic journals, and (3) promote using online platforms. Also, over the next 2 years, periodic studies will be conducted to evaluate the treatment status of mechanical ocular trauma.

## 3 | RESULTS


The guideline development group investigated patients' wishes and organized expert consensus meetings. Finally, 16 recommendations were formed. The flowchart of treatment for eyes with NLP induced by mechanical ocular trauma is shown in Figure 1.

### 3.1 | Clinical Recommendation 1: Identification of NLP eyes induced by mechanical ocular trauma

History of ocular trauma and visual acuity examinations are critical in the identification of NLP (GPS).

#### 3.1.1 | Recommendation explanations

NLP eyes should be examined in a dark room, and the other eye should be completely covered. The light from the direct or



**FIGURE 1** Flowchart of treatment for eyes with no light perception induced by mechanical ocular trauma

indirect ophthalmoscope, candle, and flashlight should be used to examine nine directions including normotopia, supranasal, superior, supratemporal, temporal, inferotemporal, inferior, inferonasal, and nasal parts. Lack of response across all sections would classify as NLP.

### 3.2 | Clinical Recommendation 2: Common causes of NLP induced by mechanical ocular trauma

Common causes of traumatized eyes with NLP include the following: (1) Direct or indirect damage to the optic nerve can cause NLP in open globe injuries, and optic nerve contusion is the leading cause of closed globe injuries. (2) Extensive ocular tissue loss or severe damage to the ocular tissue (GPS).

### 3.2.1 | Recommendation explanations

Loss of visual acuity after mechanical ocular trauma is due to a combination of many factors, including media opacity caused by severe corneal injury, traumatic cataracts, dense intraocular haemorrhage, and suppurative endophthalmitis.<sup>23</sup> Other common causes of post-traumatic NLP include intraocular tissue prolapse, injury of the optic nerve, extensive damage to the retina or choroid, severe ciliary body damage, and psychological factors (such as hysteria).<sup>24</sup>

### 3.3 | Clinical Recommendation 3: Prognostic risk factors for NLP eyes induced by mechanical ocular trauma

The risk factors for poor prognosis in NLP eyes induced by mechanical ocular trauma include rupture (2D), intraocular foreign body (IOFB) retention (2C), Zone III injury (2C), wound length  $\geq 10$  mm (2C), prolapse of the retina, choroid, and vitreous (2C), the timing of primary repair  $> 24$  h (2C), excessive/overlong interval to secondary surgery (2D), lens damage (2D), severe ciliary body damage (2C), severe intraocular haemorrhage (2C), proliferative vitreoretinopathy (PVR) (2B), retinal and/or choroidal detachment (RD) (2B), relative afferent pupillary defect (2C), and traumatic endophthalmitis (2D).

### 3.3.1 | Recommendation explanations

The guideline development group determined the outcome indicator according to ocular trauma score variables<sup>25,26</sup> and provided a systematic review showing that eyeball rupture was the most important risk factor for poor prognosis. The incidence of low vision (Risk difference, RD = 0.75, 95% CI:0.57–0.92),<sup>27</sup> NLP ( $p = 0.15$ , 95% CI:0.07–0.30),<sup>28,29</sup> and PVR after vitrectomy is higher in ruptured eyes. The incidence of endophthalmitis significantly increases in the presence of IOFBs retention (Odds ratio, OR = 2.06, 95% CI:1.47–2.88).<sup>30,31</sup> The incidence of poor visual acuity prognosis and the possibility of long-term NLP were highest in patients with Zone III injury (OR = 5.45, 95% CI:2.65–11.21).<sup>32</sup> Comparatively, the possibility of the above two incidences of Zone I injuries was the lowest, and the incidence of RD and enucleation after Zone III injuries was the highest. If the length of the wound was greater than 10 mm, RD was most likely to occur (OR = 11.46, 95% CI:4.41–29.74).<sup>33</sup>

The timing of primary surgery and vitrectomy has a significant impact on the prognosis of mechanical ocular trauma, and a delay of primary surgery over 24 h increases the risk of endophthalmitis.<sup>34,35</sup> The risk of poor visual outcomes and RD was lower in patients who underwent vitrectomy 7–14 days after injury than in those who underwent vitrectomy  $> 14$  days after injury.<sup>36</sup> Lens injuries caused by ocular trauma can increase the incidence of poor visual prognosis and endophthalmitis. Severe ciliary body injury and intraocular haemorrhage can also affect the prognosis of visual recovery and lead to abnormal intraocular pressure (IOP).<sup>33</sup>





sterilized air bubble or viscoelastic agent after suturing, laying a good foundation for vitrectomy. For perforating wounds, anterior wounds should be sutured first, whereas posterior wounds are generally not sutured. IOFBs should be removed as soon as possible.<sup>54</sup> Traumatic cataracts are recommended to be removed during vitrectomy.

### 3.7 | Clinical Recommendation 7: Preparation for vitrectomy in NLP eyes induced by mechanical ocular trauma

Preparation for vitrectomy in NLP eyes induced by mechanical ocular trauma (including local and systemic preparation, preventive medication, requirements for surgeons, equipment and material preparation, and informed consent). (1) NLP eyes induced by mechanical ocular trauma should undergo detailed examinations before vitrectomy, and preoperative anesthesia evaluation should be performed when general anesthesia is selected. Patients and their family members should be informed of operative risks and potential postoperative complications before surgery (GPS). (2) Comprehensive surgical equipment and instruments should be prepared. The operation should be performed by experienced doctors in the field of ocular trauma or vitreoretinal surgery (GPS).

#### 3.7.1 | Recommendation explanations

Patients with NLP should undergo detailed examinations before vitrectomy for personalized surgical planning and prediction of the postoperative anatomy and visual recovery. General anesthesia should be prioritized after the patients' general conditions are assessed. Antibiotics should be used to prevent endophthalmitis. The use of antibiotics is the same as that in Clinical Recommendation 5. For patients with high IOP after primary surgery, topical eye drops, oral carbonic anhydrase inhibitors, or mannitol can be administered if necessary. Vitrectomy should be performed by an experienced ophthalmologist specializing in ocular trauma. The equipment preparation is the same as that in Clinical Recommendation 6.

### 3.8 | Clinical Recommendation 8: Indications, timing, and procedures of vitrectomy in NLP eyes induced by mechanical ocular trauma

Indications, timing, and procedures of vitrectomy in NLP eyes induced by mechanical ocular trauma. (1) The main indications for vitrectomy include vitreous haemorrhage (VH), RD, choroidal detachment, IOFB, and endophthalmitis (GPS). (2) The appropriate timing for vitrectomy should be determined according to the type and severity of ocular trauma. Most surgeons choose to perform vitrectomy 3–14 days after injury. Patients with IOFB or a high risk of infection should be immediately operated. Vitrectomy is not recommended to be performed later than 14 days after ocular trauma for the cause of severe PVR (GPS). (3)

Routine surgical procedures include anterior segment reconstruction and reattachment of retina and choroid (GPS).

#### 3.8.1 | Recommendation explanations

There is a view that RD and repeated VH during conservative treatment after blunt trauma are indications for vitrectomy. Currently, the timing is mainly divided into early surgery ( $\leq 3$  days), delayed surgery (4–7 days), late surgery (8–14 days), and very late surgery ( $> 14$  days).<sup>55</sup> Vitrectomy is usually performed as early as possible to reduce the incidence of endophthalmitis in patients with IOFBs or a high risk of infection. Early vitrectomy has a certain effect on preventing proliferation. However, the indication is constricted, such as without corneal rupture and large ocular wound. Suprachoroidal haemorrhage should be drained during vitrectomy, and a temporary corneal prosthesis or endoscope can be used in patients with corneal blood staining. If vitrectomy cannot be conducted easily, bimanual manipulation can be adopted with chandelier illumination. If the choroidal incarceration is in the scleral wound or if there is a large range of choroid avulsion, the choroid can be sutured after cleaning the choroidal lesion and fixed on the sclera.

### 3.9 | Clinical Recommendation 9: Prevention and treatment of intraoperative complications during vitrectomy for NLP eyes

Prevention and treatment of intraoperative complications during vitrectomy for NLP eyes. (1) Intraoperative haemorrhage. When removing IOFBs from the ocular wall, photocoagulation or cautery should be performed on the retina around the foreign body before removal. Sufficient cautery should be performed at the appropriate site before retinotomy or retinectomy to prevent bleeding. In cases of intraoperative bleeding, an increased infusion pressure or cautery should be conducted (GPS). (2) Iatrogenic retinal tears: Iatrogenic retinal tears can be prevented by careful manipulation. The present retinal tears can be closed by laser photocoagulation or cryotherapy, and silicone oil or inert gas is filled in the eye at the end of vitrectomy (GPS). (3) Perfluorocarbon liquid migrates into the subretinal space. When the perfluorocarbon liquid migrated into the subretinal space, it can be removed with a back-flush needle. Then the proliferative membrane peeling, retinotomy, or retinectomy were performed followed by silicone oil filling (GPS). (4) Silicone oil goes under the retina. When silicone oil goes under the retina, it can be removed by retinectomy or retinotomy at the appropriate site (GPS).

#### 3.9.1 | Recommendation explanations

Intraoperative bleeding is mainly caused by inadequate cautery or accidental damage to the retinal vessels during the separation of the proliferative membrane adhered to the retina or near the wound. If there is intraoperative bleeding, hemostasis can be achieved by

increasing infusion pressure or cautery. If large and persistent bleeding results in blurred media and the bleeding point cannot be accurately identified, the air-fluid exchange can be temporarily performed, and intraocular cautery should be used to stop the bleeding after the intraoperative field is clear.

Possible causes of intraoperative iatrogenic retinal tears include vitreous traction when surgical instruments repeatedly pass in and out of the sclerotomy and peeling of the epiretinal membrane. According to the results of the meta-analysis, the incidence of iatrogenic retinal tears in vitrectomy is 11%, which can be reduced by being careful during the operation, selecting proper surgical instruments, and reducing the frequency of intraocular instruments in and out.<sup>56</sup> For present iatrogenic retinal tears, laser photocoagulation or cryotherapy can be used, and silicone oil or C3F8 can be filled in the eyes at the end of vitrectomy.

Perfluorocarbon liquid that migrates into the subretinal space is mainly caused by inadequate removal of epi-retinal proliferation or incomplete retinectomy or retinotomy. Perfluorocarbon liquids should be removed, and retinal lesions should be treated thoroughly.

The silicone oil migrated to the subretinal space mainly because the silicone oil injection needle did not directly enter the vitreous cavity, or the edges of the large retinal tears were not completely attached. Silicone oil can be injected after the retinal lesions have been completely cleared.

### 3.10 | Clinical Recommendation 10: Prevention and treatment of complications after vitrectomy in NLP eyes induced by mechanical ocular trauma

Prevention and treatment of complications after vitrectomy in NLP eyes induced by mechanical ocular trauma. (1) Corneal edema: Corneal edema usually gradually recovers after local treatment (GPS). (2) Hyphema: Iris damage should be avoided during vitrectomy to prevent postoperative hyphema. If the amount of hyphema is large and IOP is high, hyphema removal should be conducted as early as possible, and hemostatic drugs should be administered (GPS). (3) Complicated cataracts caused by silicone oil tamponade. Patients should be treated with cataract extraction combined with oil removal (GPS). (4) Vitreous or retinal haemorrhage. Bleeding can be prevented by thorough hemostasis during vitrectomy. Reoperation is required in patients with recurrent, unabsorbed, and severe VH (GPS). (5) Recurrent retinal or choroidal detachment: The iatrogenic retinal injury should be avoided, and retinal and choroidal lesions should be treated thoroughly. Further surgery should be considered prudently if there is no chance to recover vision (GPS). (6) Silicone oil-dependent eyes. Delayed removal of silicone oil or reoperation to replace it (GPS). (7) Abnormal IOP. There is currently no effective treatment for low IOP caused by severe ciliary body damage. Temporarily high IOP should be treated according to different causes. Anti-glaucoma surgery is required if the drug cannot control secondary glaucoma (GPS). (8) Traumatic endophthalmitis can be rescued by vitrectomy combined with silicone oil tamponade and intravitreal antibiotics (GPS).

### 3.10.1 | Recommendation explanations

Meta-analysis indicated that 31% of patients showed varying degrees of corneal edema after vitrectomy, caused by corneal injury, early operation time, long operation duration, or post-traumatic intraocular inflammation, usually recovers gradually after local treatment.<sup>57,58</sup> The incidence of postoperative hyphema is 10%. Treatment of postoperative hyphema is based on the amount of bleeding. If the amount of hyphema is large and IOP is high, hyphema should be removed in time.<sup>59</sup> Approximately, 16% of patients develop complicated cataracts. Vitrectomy combined with cataract surgery should be performed for complicated cataracts caused by secondary glaucoma or endophthalmitis. Complicated cataracts caused by silicone oil tamponade should be treated with silicone oil removal and cataract extraction.

The main reasons for postoperative VH may be the early vitrectomy time, accidental injury during operation, or incomplete hemostasis of intraoperative retinal bleeding. A small amount of retinal haemorrhage only needs to be followed up without any treatment. If vitreous and retinal bleeding is severe, further surgery may be required. The incidence of postoperative RD is about 10%, caused by the recurrent PVR,<sup>60</sup> and reoperation should be considered.

Seven per cent of patients need to delay the removal time of silicone oil, even leading to silicone oil dependence. A postponed removal of silicone oil could be considered in patients with intended RD after silicone oil removal or persistently low IOP, and long-term silicone oil tamponade can effectively maintain a certain IOP as shown in the results of the meta-analysis.<sup>61,62</sup> Silicone oil may be emulsified owing to its long-term retention in 8% of patients and should be removed accordingly. In silicone oil-dependent eyes, emulsified silicone oil can be exchanged with new silicone oil. Low IOP can be caused by severe ciliary body damage, retinal or choroidal defects, or PVR in 10% of patients.<sup>63,64</sup> Temporary high IOP in the early postoperative period could be caused by excessive silicone oil tamponade, inflammatory reactions in the anterior chamber, or closure of the peripheral iridectomy. Secondary glaucoma occurs in 11% of patients because of postoperative silicone oil emulsification, hyphema, or anterior chamber angle damage induced by blunt trauma. Anti-glaucoma surgery is required if the drug cannot control secondary glaucoma.<sup>23,65–67</sup>

Patients with traumatic endophthalmitis should be administered intravitreal antibiotics during surgery.<sup>68–72</sup> However, approximately 4% of patients still end up with eyeball atrophy, mostly caused by ciliary body damage induced by severe mechanical ocular trauma, recurrence of RD, or incomplete treatment of endophthalmitis.<sup>73–76</sup>

### 3.11 | Clinical Recommendation 11: Choice of intraocular tamponade in vitrectomy for NLP eyes

Silicone oil is recommended as the main intraocular tamponade for NLP eyes. Inert gas or balanced salt solution (BSS) can be used for minor retinal injuries. (1D)



### 3.11.1 | Recommendation explanations

Common tamponades in vitrectomy include silicone oil, inert gas, and BSS. Inert gas tamponade can promote retinal reattachment and absorption of subretinal fluid, which is suitable for medium- and short-term intraocular tamponades. BSS has the highest histocompatibility but does not have an expansion effect on the retina. Silicone oil has a longer duration owing to its lower density and higher surface tension than water.<sup>77</sup> Additionally, studies have shown that silicone oil can inhibit the growth of residual pathogenic bacteria in the eye.<sup>78,79</sup>

### 3.12 | Clinical Recommendation 12: Decisions on either salvage of the globe or enucleation in NLP eyes induced by mechanical ocular trauma (including surgical indication, contraindication, and medical ethics)

Decisions on either salvage of the globe or enucleation in NLP eyes induced by mechanical ocular trauma (including surgical indication, contraindication, and medical ethics). (1) Enucleation of the eyeball is not recommended in emergency surgery unless almost all intraocular tissues are lost (GPS). (2) The possibility of enucleation of traumatic eyes after the primary operation cannot be ruled out when severe complications such as sympathetic ophthalmia, long-term eye pain, and eyeball atrophy occur, and the possibility of restoring light perception is lost (GPS). (3) If enucleation is inevitable, ophthalmologists should fully communicate with patients and their family members and provide psychological counseling to relieve patients' anxiety (GPS).

#### 3.12.1 | Recommendation explanations

NLP eyes can recover vision after mechanical ocular trauma, and NLP before the primary operation cannot be used as an indication of permanent visual loss.<sup>9,33</sup> Debridement should be performed during primary surgery, and damage to the intraocular tissue should be evaluated carefully at the same time. Vitrectomy should be performed on time in patients with potential visual recovery, and enucleation should be judged again during vitrectomy. Enucleation can be considered in the primary operation under the following conditions: (1) The affected eye has NLP before ocular trauma, meanwhile, almost all intraocular content is lost, and the anatomical structure of the eyeball is seriously damaged. (2) The anatomical structure of the eyeball is severely damaged and cannot undergo vitrectomy or any other subsequent treatment in aged NLP patients with severe systemic diseases. (3) Almost all intraocular content loss after trauma and the eyeball cannot be repaired. Ophthalmologists should be aware of NLP caused by psychosocial and mental factors before deciding on enucleation and assess whether there is any possibility of vision recovery.<sup>80,81</sup> Enucleation may cause severe psychological and emotional trauma to patients;

hence, ophthalmologists should provide psychological counseling and humanistic care to relieve patients' anxiety.

### 3.13 | Clinical Recommendation 13: Criteria and prognosis of silicone oil-dependent eyes after vitrectomy for NLP induced by mechanical ocular trauma

Criteria and prognosis of silicone oil-dependent eyes after vitrectomy for NLP induced by mechanical ocular trauma. (1) The criteria for silicone oil-dependent eyes are limited potential for visual recovery or recurrent RD after silicone oil removal, persistent low IOP, and functionally monocular patients (1C). (2) Long-term silicone oil tamponade effectively maintains certain IOP (1D).

#### 3.13.1 | Recommendation explanations

There are several problems with silicone oil as a long-term intraocular tamponade, including silicone oil emulsification, secondary glaucoma, complicated cataracts, and corneal degeneration. Silicone oil should be removed 3–6 months after vitrectomy.<sup>82</sup> However, in patients with severe injury, recurrent RD and persistently low IOP may occur if silicone oil is removed, leading to eyeball atrophy and potential enucleation. Hence, some patients need to extend the silicone oil removal time, or even need to permanently retain silicone oil and become silicone oil-dependent eyes. Meta-analysis showed that if NLP eyes have experienced several complications, such as the limited recovery of visual potential, recurrent RD after silicone oil removal, persistent low IOP, or functionally monocular patients, persistent silicone oil tamponade could be considered.<sup>61,62</sup> Long-term silicone oil tamponade can effectively maintain certain IOP and has no obvious effect on vision improvement or retinal reattachment.<sup>23,83</sup>

### 3.14 | Clinical Recommendation 14: Follow-up precautions after vitrectomy for NLP eyes induced by mechanical ocular trauma

Routine follow-up should be performed within 1 week and at 1, 3, 6, and 12 months after vitrectomy. Visual acuity, IOP, and ocular structure should be examined during follow-up. Complications, such as endophthalmitis, PVR, silicone oil emulsification, keratopathy, secondary glaucoma, and sympathetic ophthalmia should be treated on time (GPS).

#### 3.14.1 | Recommendation explanations

Follow-up was performed no less than 6 months after vitrectomy. In most studies, the longest follow-up was 2–3 years.<sup>23,57,84–87</sup>

### 3.15 | Clinical recommendation 15. Visual outcomes and recovery process after vitrectomy for NLP eyes induced by mechanical ocular trauma

The final visual acuity of NLP eyes induced by mechanical ocular trauma has the potential to improve after vitrectomy. The rate of vision recovery gradually increases after vitrectomy (2D).

#### 3.15.1 | Recommendation explanations

Meta-analysis showed that visual acuity was restored to light perception and hand motion in 34% of eyes with NLP after vitrectomy,<sup>8,9,23,42,85,86,88</sup> recovered from NLP to counting fingers (CF)-20/400 in 16% and had a long-term visual outcome better than 20/400 in 11%. Subgroup analysis showed that the light perception recovery rate gradually increased until 1 year after vitrectomy.

### 3.16 | Clinical Recommendation 16: Effectiveness of corticosteroids and antibiotics in the treatment of NLP induced by mechanical ocular trauma

Effectiveness of corticosteroids and antibiotics in the treatment of NLP induced by mechanical ocular trauma. (1) For patients with NLP eyes induced by mechanical ocular trauma, the use of triamcinolone acetonide or dexamethasone after primary surgery is recommended as single intravitreal drug therapy for empirical anti-inflammatory therapy (2D). (2) For patients with NLP induced by mechanical ocular trauma, vancomycin is recommended as a single intravitreal drug empirical anti-infective therapy. However, concomitant medicine with vancomycin is also recommended based on the outcomes of pathogen culture and antimicrobial susceptibility tests, which include ceftazidime, ceftriaxone, cefaloridine, cefuroxime, cefepime, moxifloxacin, levofloxacin, ofloxacin, clindamycin, and tobramycin. Voriconazole and amphotericin have been recommended for the treatment of fungal infections (2D).

#### 3.16.1 | Recommendation explanations

Meta-analysis showed that intravitreal injection of triamcinolone acetonide can effectively reduce the incidence of postoperative ocular inflammation during complex ocular trauma surgery. Intraocular injection of vancomycin can effectively improve the surgical effect in mechanical ocular trauma patients with NLP.<sup>89-91</sup> In patients with penetrating ocular injuries and IOFBs, a single intravitreal injection of vancomycin can help reduce the incidence of endophthalmitis during the treatment process (RR = 0.24, 95% CI:0.09-0.63,  $p < 0.05$ ).<sup>92,93</sup> Vancomycin combined with ceftazidime was used for the treatment of open globe injury, and systemic or intraocular injection can reduce the incidence of post-traumatic endophthalmitis and the rate of enucleation, and improve the prognosis

of infectious endophthalmitis. Combined vitreous injection of vancomycin and ceftazidime during surgical treatment has a certain effect on postoperative visual acuity improvement in patients with open globe injury and endophthalmitis. Intraocular injection of vancomycin and ceftazidime, combined with systemic administration of cefuroxime, cefazolin, or clindamycin, can reduce the rate of enucleation after trauma.<sup>94-96</sup> Systemic intravenous vancomycin combined with cefepime, tobramycin, or ofloxacin can reduce the incidence of post-traumatic endophthalmitis.<sup>97</sup> It has been reported that hypersensitivity caused by intravitreal vancomycin may lead to haemorrhagic obstructive retinal vasculitis (HORV). Regular postoperative follow-up for ocular fundus examination should be performed. If HORV is suspected, further intraocular injection of vancomycin should be avoided.<sup>98,99</sup>

## 4 | CONCLUSION

This guideline focuses on the diagnosis, treatment, and prognosis of mechanical ocular trauma with NLP in strict accordance with WHO guideline formulation specifications and international standards. The guideline development group surveyed ophthalmologists and experts in the ocular trauma field, collected and arranged key questions on mechanical ocular trauma with NLP, comprehensively retrieved and evaluated systemically relevant research evidence, conducted in-depth interviews, and analyzed patient preferences and values. Eventually, 16 recommendations directed at the diagnosis, treatment, and prognosis of mechanical ocular trauma with NLP were developed. Each recommendation has been explained in detail, providing important guidance for the diagnosis, treatment, and long-term prognosis prediction of mechanical ocular trauma with NLP. The guideline development group reminds us that the level of some recommendations is weak because of the inadequacy or lack of clinical evidence. Ophthalmologists should receive relevant training when using this guideline. Valuable suggestions and opinions from users on the shortcomings of this guideline are warmly welcomed to ensure continuous improvement. In addition, this guideline is not intended for commercial promotion or publicity purposes.

## CONFLICT OF INTEREST

The authors declare no conflict of interest.

## ORCID

Hua Yan  <https://orcid.org/0000-0002-7651-840X>

Kehu Yang  <https://orcid.org/0000-0001-7864-3012>

Andres Rousselot  <https://orcid.org/0000-0003-2341-0988>

Shihui Wei  <https://orcid.org/0000-0003-2910-0446>

Yaolong Chen  <https://orcid.org/0000-0002-7338-4418>

## REFERENCES

1. Pieramici DJ, Sternberg P, Aaberg TM, et al. A system for classifying mechanical injuries of the eye (globe). The Ocular Trauma Classification Group. *Am J Ophthalmol*. 1997;123:820-831.

2. Liu Y, Feng K, Jiang H, et al. Characteristics and treatments of ocular blast injury in Tianjin explosion in China. *BMC Ophthalmol*. 2020;20(1):185.
3. Fujikawa A, Mohamed YH, Kinoshita H, et al. Visual outcomes and prognostic factors in open-globe injuries. *BMC Ophthalmol*. 2018;18(1):138.
4. Desai P, Morris DS, Minassian DC, Macewen CJ. Trends in serious ocular trauma in Scotland. *Eye (London)*. 2105;29(5):611-618.
5. China Ocular Trauma Society of Chinese Ophthalmological Society: Consensus of Chinese experts on the treatment of mechanical ocular trauma with no light perception (2020). *Chin J Ophthalmol*. 2020;56:815-819.
6. Dong F, Dai R. The surgical treatment of traumatized eyes with no light perception. *Zhonghua Yan Ke Za Zhi*. 2002;38:657-659.
7. Salehi-Had H, Andreoli CM, Andreoli MT, Kloek CE, Mukai S. Visual outcomes of vitreoretinal surgery in eyes with severe open-globe injury presenting with no-light-perception vision. *Graefes Arch Clin Exp Ophthalmol*. 2002;247:477-483.
8. Feng K, Hu YT, Ma Z. Prognostic indicators for no light perception after open-globe injury: eye injury vitrectomy study. *Am J Ophthalmol*. 2011;152:654-662.
9. Agrawal R, Wei HS, Teoh S. Predictive factors for final outcome of severely traumatized eyes with no light perception. *BMC Ophthalmol*. 2012;12:16.
10. Burda BU, Chambers AR, Johnson JC. Appraisal of guideline developed by the World Health Organization. *Public Health*. 2014;128:444-474.
11. Graham R, Mancher M, Miller Wolman D, Greenfield S, Steinberg E, eds. *Clinical Practice Guideline We Can Trust*. National Academies Press (US); 2011.
12. Brouwers MC, Kho ME, Browman GP, et al. AGREE II: advancing guideline development, reporting and evaluation in health care. *Cmaj*. 2010;182:E839-E842.
13. Chen Y, Yang K, Marušić A, et al. A reporting tool for practice guideline in health care: the right statement. *Ann Intern Med*. 2017;166:128-132.
14. Li X, Wei L, Shang W, et al. Trace and evaluation systems for health services quality in rural and remote areas: a systematic review. *J Public Health*. 2018;26:127-135.
15. Pan B, Ge L, Xun Y-Q, et al. Exercise training modalities in patients with type 2 diabetes mellitus: a systematic review and network meta-analysis. *Int J Behav Nutr Phys Act*. 2018;15:72.
16. Shea BJ, Reeves BC, Wells G, et al. AMSTAR 2: a critical appraisal tool for systematic reviews that include randomised or non-randomised studies of healthcare interventions, or both. *BMJ*. 2017;358:j4008.
17. Wells GA, Shea BJ, O'Connell D, et al. The Newcastle-Ottawa Scale (NOS) for assessing the quality of non-randomised studies in meta-analysis. 2013. [http://www.ohri.ca/programs/clinical\\_epidemiology/oxford.asp](http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp)
18. Higgins JPT, Altman DG, Gotzsche PC, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ*. 2011;343:d5928.
19. Whiting PF. QUADAS-2: a revised tool for the quality assessment of diagnostic accuracy studies. *Ann Intern Med*. 2011;155:529-536.
20. Jaeschke R, Guyatt GH, Dellinger P, et al. Use of GRADE grid to reach decisions on clinical practice guideline when consensus is elusive. *BMJ*. 2008;337:a744.
21. Norris SL, Meerpohl JJ, Akl EA, et al. The skills and experience of GRADE methodologists can be assessed with a simple tool. *J Clin Epidemiol*. 2016;79:150-158.
22. Guyatt GH, Alonso-Coello P, Schünemann HJ, et al. Guideline panels should seldom make good practice statements: guidance from the GRADE Working Group. *J Clin Epidemiol*. 2016;80:3-7.
23. Yang SS, Jiang T. Vitrectomy combined with silicone oil tamponade in the treatment of severely traumatized eyes with the visual acuity of no light perception. *Int J Ophthalmol*. 2013;6:198-203.
24. Agrawal R, Shah M, Mireskandari K, Yong GK. Controversies in ocular trauma classification and management: review. *Int Ophthalmol*. 2013;33:435-445.
25. Kuhn F, Maisiak R, Mann L, Mester V, Morris R, Witherspoon C. The Ocular Trauma Score (OTS). *Ophthalmol Clin North Am*. 2002;15(2):163-165.
26. Brundridge W, Reed D, Santamaria J, Mehta A, Valentin F, Davies B. Open globe trauma in a military hospital: a review of the Ocular Trauma Score to help predict enucleation or evisceration. *Graefes Arch Clin Exp Ophthalmol*. 2019;257(8):1789-1793.
27. Supreeyathitikul P, Chokesuwattanasul S, Choovuthayakorn J, et al. Epidemiology and outcomes following open globe injury in agricultural region, an 11-year experience. *Ophthalmic Epidemiol*. 2020;27:246-251.
28. Soheilian M, Peyman GA, Wafapoor H, Navarro GC, Thompson H. Surgical management of traumatic retinal detachment with perfluorocarbon liquid. The Vitreous Study Group. *Int Ophthalmol*. 1996;20:241-249.
29. Shrestha SM, Anthony CL, Justin GA, et al. Factors affecting final functional outcomes in open-globe injuries and use of ocular trauma score as a predictive tool in Nepalese population. *BMC Ophthalmol*. 2021;21:69.
30. Andreoli CM, Andreoli MT, Kloek CE, Ahuero AE, Vavvas D, Durand ML. Low rate of endophthalmitis in a large series of open globe injuries. *Am J Ophthalmol*. 2009;147:601-608.
31. Asencio MA, Huertas M, Carranza R, Tenias JM, Celis J, Gonzalez-Del Valle F. A case-control study of post-traumatic endophthalmitis at a Spanish hospital. *Int Ophthalmol*. 2016;36:185-194.
32. Yalcin Tök O, Tok L, Eraslan E, Ozkaya D, Ornek F, Bardak Y. Prognostic factors influencing final visual acuity in open globe injuries. *J Trauma*. 2011;71:1794-1800.
33. Feng K, Shen L, Pang X, et al. Case-control study of risk factors for no light perception after open-globe injury: eye injury vitrectomy study. *Retina*. 2011;31:1988-1996.
34. Guven S, Durukan AH, Erdurman C, Kucukcilioglu M. Prognostic factors for open-globe injuries: variables for poor visual outcome. *Eye (London, England)*. 2019;33(3):392-397.
35. Jovanovic N, Peek-Asa C, Young T, Nisic F, Musa S. Eye injury and demographic parameters associated with poor visual outcome. *J Fr Ophthalmol*. 2019;42(8):864-873.
36. Bekele S, Gelaw Y. Pattern and prognostic factor of ocular injuries in southwest Ethiopia: a hospital based prospective study. *Int Eye Sci*. 2016;16(5):811-817.
37. Thompson WS, Rubsamen PE, Flynn HW, Schiffman J, Cousins SW. Endophthalmitis after penetrating trauma. Risk factors and visual acuity outcomes. *Ophthalmology*. 1995;102:1696-1701.
38. Lieb DF, Scott IU, Flynn HW, Miller D, Feuer WJ. Open globe injuries with positive intraocular cultures: factors influencing final visual acuity outcomes. *Ophthalmology*. 2003;110:1560-1566.
39. Essex R, Yi Q, Charles P, Allen P. Post-traumatic endophthalmitis. *Ophthalmology*. 2004;111:2015-2022.
40. Schmidt GW, Broman AT, Hindman HB, Grant MP. Vision survival after open globe injury predicted by classification and regression tree analysis. *Ophthalmology*. 2008;115:202-209.
41. Al-Mezaine HS, Osman EA, Kangave D, Abu El-Asrar AM, Abu El-Asrar AM. Risk factors for culture-positive endophthalmitis after repair of open globe injuries. *Eur J Ophthalmol*. 2020;20:201-208.
42. Soni NG, Bauza AM, Son JH, Langer PD, Zarbin MA, Bhagat N. Open globe ocular trauma: functional outcome of eyes with no light perception at initial presentation. *Retina*. 2013;33:380-386.
43. Zhang Y, Zhang M, Jiang C, Qiu HY. Intraocular foreign bodies in China: clinical characteristics, prognostic factors, and visual outcomes in 1,421 eyes. *Am J Ophthalmol*. 2011;152:66-73.

44. Duan F, Yuan Z, Liao J, Zheng Y, Yang Y, Lin X. Incidence and risk factors of intraocular foreign body-related endophthalmitis in Southern China. *J Ophthalmol*. 2018;2018:8959108.
45. Gan YF, Cao JR, Song GW, Zheng HH, Dong ZZ. Analysis of the visual outcomes and clinical risk factors for the occurrence of intraocular foreign-body related endophthalmitis. *Chin J Optom Ophthalmol Vis Sci*. 2021;23:109-115.
46. Ahmed Y, Schimel AM, Pathengay A, Colyer MH, Flynn HW. Endophthalmitis following open-globe injuries. *Eye (London)*. 2012;26(2):212-217.
47. Huang JM, Pansick AD, Blomquist PH. Use of intravenous vancomycin and cefepime in preventing endophthalmitis after open globe injury. *J Ocul Pharmacol Ther*. 2016;32(7):437-441.
48. Silpa-Archa S, Kumsiang K, Chotcomwongse P, Preble JM, Stephen Foster C. Poor prognostic factors in post-traumatic endophthalmitis following open globe injury. *Int J Ophthalmol*. 2020;13(12):1968-1975.
49. Abouammoh MA, Al-Mousa A, Gogandi M, et al. Prophylactic intravitreal antibiotics reduce the risk of post-traumatic endophthalmitis after repair of open globe injuries. *Acta Ophthalmol*. 2018;96(3):e361-e365.
50. Thompson JT, Parver LM, Enger CL, Mieler WF, Liggett PE. Infectious endophthalmitis after penetrating injuries with retained intraocular foreign bodies. National Eye Trauma System. *Ophthalmology*. 1993;100:1468-1474.
51. Blanch RJ, Bishop J, Javidi H, Murray PI, Murray PI. Effect of time to primary repair on final visual outcome after open globe injury. *Br J Ophthalmol*. 2019;103:1491-1494.
52. Scott IU, McCabe CM, Flynn HW, et al. Local anesthesia with intravenous sedation for surgical repair of selected open globe injuries. *Am J Ophthalmol*. 2002;134:707-711.
53. McClellan AJ, Daubert JJ, Relhan N, Tran KD, Flynn HW, Gayer S. Comparison of regional vs. general anesthesia for surgical repair of open-globe injuries at a University Referral Center. *Ophthalmol Retina*. 2017;1:188-191.
54. Yan H. The importance of the primary surgical management of open globe injury. *Chin J Optom Ophthalmol Vis Sci*. 2015;17(2):65-67.
55. Kuhn F. The timing of reconstruction in severe mechanical trauma. *Ophthalmic Res*. 2014;51:67-72.
56. Batman C, Çekiç O, Totan Y, Ozkan SS, Zilelioglu O. Combined phacoemulsification, vitrectomy, foreign-body extraction, and intraocular lens implantation. *J Cataract Refract Surg*. 2000;26(2):254-259.
57. Pimolrat W, Choovuthayakorn J, Watanachai N, et al. Predictive factors of open globe injury in patients requiring vitrectomy. *Injury*. 2014;45:212-216.
58. Yu H, Li J, Yu Y, et al. Optimal timing of vitrectomy for severe mechanical ocular trauma: a retrospective observational study. *Sci Rep*. 2019;9:18016.
59. Singh R, Bhalekar S, Dogra M, Gupta A. 23-gauge vitrectomy with intraocular foreign body removal via the limbus: an alternative approach for select cases. *Indian J Ophthalmol*. 2014;62:707-710.
60. Wang R, Wang XL, Wang Y, et al. Endotamponades in pars plana vitrectomy for metallic intraocular foreign body associated with endophthalmitis. *Int J Ophthalmol*. 2011;4:95-99.
61. Stappeler T, Morphis G, Irigoyen C, Heimann H. Is there a role for long-term silicone oil tamponade for more than twelve months in vitreoretinal surgery? *Ophthalmologica*. 2011;226(1):36-41.
62. Vidne-Hay O, Platner E, Alhalel A, Moisseiev J. Long-term silicone oil tamponade in eyes with complicated retinal detachment. *Eur J Ophthalmol*. 2021;2021:11206721211019551.
63. Wang YN, Shen LJ, Wang CG, et al. Prognosis of traumatic eyes with no light perception undergone vitrectomy and analysis of risk factors. *Chin J Ophthalmol*. 2007;43:340-345.
64. Xu ZJ, Xu YM, Lin HR, Lin S, Qin B. Vitrectomy for severe ocular injury with no light perception. *Int J Ophthalmol*. 2010;10:1370-1371.
65. Girkin CA, McGwin G, Long C, Morris R, Kuhn F. Glaucoma after ocular contusion: a cohort study of the United States Eye Injury Registry. *J Glaucoma*. 2005;14:470-473.
66. Szurman P, Roters S, Grisanti S, et al. Primary silicone oil tamponade in the management of severe intraocular foreign body injuries: an 8-year follow-up. *Retina*. 2007;27:304-311.
67. Bai H-Q, Yao L, Wang D-B, Jin R, Wang Y-X. Causes and treatments of traumatic secondary glaucoma. *Eur J Ophthalmol*. 2009;19:201-206.
68. Mieler WF, Ellis MK, Williams DF, Han DP. Retained intraocular foreign bodies and endophthalmitis. *Ophthalmology*. 1990;97:1532-1538.
69. Abu El-Asrar AM, Al-Amro SA, Khan NM, Kangave D. Retinal detachment after posterior segment intraocular foreign body injuries. *Int Ophthalmol*. 1998;22:369-375.
70. Jonas JB, Knorr HLJ, Budde WM. Prognostic factors in ocular injuries caused by intraocular or retrobulbar foreign bodies. *Ophthalmology*. 2000;107:823-828.
71. Szijsztó Z, Gaál V, Kovács B, Kuhn F. Prognosis of penetrating eye injuries with posterior segment intraocular foreign body. *Graefes Arch Clin Exp Ophthalmol*. 2008;246:161-165.
72. Sborgia G, Recchimirzo N, Niro A, Sborgia L, Sborgia A, Alessio G. 25-Gauge vitrectomy in open eye injury with retained foreign body. *J Ophthalmol*. 2017;2017:3161680.
73. Bartz-Schmidt KU, Bermig J, Kirchhof B, Wiedemann P, Walter P, Heimann K. Prognostic factors associated with the visual outcome after vitrectomy for endophthalmitis. *Graefes Arch Clin Exp Ophthalmol*. 1996;234(S1):S51-S58.
74. Kaynak S, Oner FH, Koçak N, Cingil G. Surgical management of postoperative endophthalmitis: comparison of 2 techniques. *J Cataract Refract Surg*. 2003;29:966-969.
75. Yan H, Lu Y, Yu J, et al. Silicone oil in the surgical treatment of traumatic endophthalmitis. *Eur J Ophthalmol*. 2008;18:680-684.
76. Chen X, Zha Y, Du S, Yang X. Timely use of conventional vitrectomy and endoscope-assisted vitrectomy for endophthalmitis following open ocular trauma: a retrospective study of 18 patients. *Med Sci Monit*. 2019;25:8628-8636.
77. Wagenfeld L, Zeitz O, Skevas C, Richard G. Long-lasting endotamponades in vitreoretinal surgery. *Ophthalmologica*. 2010;224:291-300.
78. Siqueira RC, Gil ADC, Canamary F, Minari M, Jorge R. Pars plana vitrectomy and silicone oil tamponade for acute endophthalmitis treatment. *Arq Bras Oftalmol*. 2009;72:28-32.
79. Yan H, Li J. Experimental study on antiviral activity of silicone oil in vitro. *Graefes Arch Clin Exp Ophthalmol*. 2008;246:1285-1289.
80. Soares IP, França VP. Evisceration and enucleation. *Semin Ophthalmol*. 2010;25:94-97.
81. Boiko EV, Churashov SV, Haritonova NN, Budko AA. Vitreoretinal surgery in the management of war-related open-globe injuries. *Graefes Arch Clin Exp Ophthalmol*. 2013;251:637-644.
82. Jonas JB, Budde WM, Knorr HLJ. Timing of retinal redetachment after removal of intraocular silicone oil tamponade. *Am J Ophthalmol*. 1999;128:628-631.
83. Feng K, Ma Z-Z. Clinical features, anatomical and visual outcomes, and prognostic factors in closed globe injuries presenting with no light perception: eye injury vitrectomy study. *Acta Ophthalmol*. 2012;90:e493-e494.
84. Warrasak S, Euswas A, Hongsakorn S. Posterior segment trauma: types of injuries, result of vitreo-retinal surgery and prophylactic broad encircling scleral buckle. *J Med Assoc Thai*. 2005;88:1916-1930.
85. Heidari E, Taheri N. Surgical treatment of severely traumatized eyes with no light perception. *Retina*. 2010;30:294-299.
86. Han Y, Kavoussi S, Adelman R. Visual recovery following open globe injury with initial no light perception. *Clin Ophthalmol*. 2015;9:1443-1448.
87. He Y, Zhang L, Wang F, Zhu M, Wang Yi, Liu Y. Timing influence on outcomes of vitrectomy for open-globe injury: a prospective randomized comparative study. *Retina*. 2020;40:725-734.



88. Bhagat N, Turbin R, Langer P, et al. Approach to management of eyes with no light perception after open globe injury. *J Ophthalmic Vis Res*. 2016;11:313-318.
89. Haider SA, Hassett P, Bron AJ. Intraocular vancomycin levels after intravitreal injection in post cataract extraction endophthalmitis. *Retina*. 2001;21:210-213.
90. Wirosko WJ, Kivlin JD. Successful treatment of orthodontic-associated traumatic endophthalmitis. *Am J Ophthalmol*. 2002;134:449-450.
91. Lipsky L, Barrett G. Intracameral antibiotics for prophylaxis of post-operative endophthalmitis in Australia: a review. *Clin Exp Ophthalmol*. 2019;47:537-541.
92. Duan YQ, Lin D, Chen ZP, Wang QC. Clinical study on the incidence of vancomycin intraocular injection in treatment with suppurative endophthalmitis. *Int Eye Sci*. 2015;15(5):917-919.
93. Shi J, Bai ZX, Tian J, Luo S, Chang HF. The prevention of infectious endophthalmitis after intraocular foreign body injury in Tibet high altitude district. *Chin J Ocul Traum Occupat Eye Dis*. 2015;37(2):112-114.
94. Miller JJ, Scott IU, Flynn HW, et al. Endophthalmitis caused by *Bacillus* species. *Am J Ophthalmol*. 2008;145:883-888.
95. Drumright RT, Regan KA, Lin AL, Moroux MG, Iyer SSR. Utility of wound cultures in the management of open globe injuries: a 5-year retrospective review. *J Ophthalmic Inflamm Infect*. 2020;10:5.
96. Mei F, Lin J, Liu M, Yang Y, Lin X, Duan F. Posttraumatic *Bacillus cereus* Endophthalmitis: clinical characteristics and antibiotic susceptibilities. *J Ophthalmol*. 2021;2021:6634179.
97. Wade Pd, Khan Ss, Khan Md. Endophthalmitis: magnitude, treatment and visual outcome in northwest frontier province of Pakistan. *Ann Afr Med*. 2009;8:19-24.
98. Witkin AJ, Chang DF, Jumper JM, et al. Vancomycin-Associated hemorrhagic occlusive retinal vasculitis: clinical characteristics of 36 eyes. *Ophthalmology*. 2017;124:583-595.
99. Oli A, Balakrishnan D. Hemorrhagic occlusive retinal vasculitis following prophylactic intraocular vancomycin injection. *JAMA Ophthalmol*. 2020;138:e195673.

**How to cite this article:** Yan H, Yang K, Ma Z, et al. Guideline for the treatment of no light perception eyes induced by mechanical ocular trauma. *J Evid Based Med*. 2022;15:302–314. <https://doi.org/10.1111/jebm.12496>